

MEDICATION GUIDE

Oxycodone (ox' i koe' done hye' droe klor' ide) Hydrochloride and Acetaminophen (a see't' a min' oh fen) Oral Solution, CII

Oxycodone Hydrochloride and Acetaminophen Oral Solution is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain, severe enough to require an opioid analgesic and for which alternative treatments are inadequate and when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

Important information about Oxycodone Hydrochloride and Acetaminophen Oral Solution:

- **Get emergency help right away if you take too much Oxycodone Hydrochloride and Acetaminophen Oral Solution (overdose).** When you first start taking Oxycodone Hydrochloride and Acetaminophen Oral Solution, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Taking Oxycodone Hydrochloride and Acetaminophen Oral Solution with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Oxycodone Hydrochloride and Acetaminophen Oral Solution. They could die from taking it. Store Oxycodone Hydrochloride and Acetaminophen Oral Solution away from children and in a safe place to prevent stealing or abuse. Selling or giving away Oxycodone Hydrochloride and Acetaminophen Oral Solution is against the law.

Do not take Oxycodone Hydrochloride and Acetaminophen Oral Solution if you have:

- Severe asthma, trouble breathing, or other lung problems.
- A bowel blockage or have narrowing of the stomach or intestines
- Known hypersensitivity to oxycodone, acetaminophen, or any ingredient in Oxycodone Hydrochloride and Acetaminophen Oral Solution

Before taking Oxycodone Hydrochloride and Acetaminophen Oral Solution, tell your healthcare provider if you have a history of:

- Head injury, seizures
- Liver, kidney, thyroid problems
- Problems urinating
- Pancreas or gallbladder problems
- Abuse of street or prescription drugs, alcohol addiction, or mental health problems

Tell your healthcare provider if you are:

- **Pregnant or planning to become pregnant.** Prolonged use of Oxycodone Hydrochloride and Acetaminophen Oral Solution during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **Breastfeeding.** Oxycodone Hydrochloride and Acetaminophen Oral Solution passes into breast milk and may harm your baby.
- Taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking Oxycodone Hydrochloride and Acetaminophen Oral Solution with certain other medicines can cause serious side effects that could lead to death.

When taking Oxycodone Hydrochloride and Acetaminophen Oral Solution:

- Do not change your dose. Take Oxycodone Hydrochloride and Acetaminophen Oral Solution exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- Ask your healthcare provider if you have any questions on how to correctly measure your dose. Always use a calibrated measuring device for Oxycodone Hydrochloride and Acetaminophen Oral Solution to correctly measure your dose. A household teaspoon or tablespoon is not an adequate measuring device. Given the inexactitude of the household spoon measure and the possibility of using a tablespoon instead of a teaspoon, which could lead to overdosage, it is strongly recommended that caregivers obtain and use a calibrated measuring device.
- Take your prescribed dose every 6 hours as needed for pain. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain.
- If you have been taking Oxycodone Hydrochloride and Acetaminophen Oral Solution regularly, do not stop taking Oxycodone Hydrochloride and Acetaminophen Oral Solution without talking to your healthcare provider.
- After you stop taking Oxycodone Hydrochloride and Acetaminophen Oral Solution, dispose of unused oral solution by flushing it down the toilet.

Continued on other side

While taking Oxycodone Hydrochloride and Acetaminophen Oral Solution DO NOT:

- Drive or operate heavy machinery, until you know how Oxycodone Hydrochloride and Acetaminophen Oral Solution affects you. Oxycodone Hydrochloride and Acetaminophen Oral Solution can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Oxycodone Hydrochloride and Acetaminophen Oral Solution may cause you to overdose and die.

The possible side effects of Oxycodone Hydrochloride and Acetaminophen Oral Solution:

- Constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help if you have:

- Trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of Oxycodone Hydrochloride and Acetaminophen Oral Solution. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to dailymed.nlm.nih.gov.**

Distributed by: FH2 Pharma LLC, Delray Beach, FL 33444

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: 09/2019

Code 1172B99